

Choose

AMNEPARA DUO™

Paracetamol Infusion IP 1.0% w/v

100 ml

Abridged Prescribing Information:

COMPOSITION: Each 100 ml contains: Paracetamol I.P, 1 g; Water for Injection I.P, Q.S. to 100 mL. **DESCRIPTION:** AMNEPARA DUO™ is clear, colorless to slightly yellowish solution in a flexible container. **INDICATIONS:** Indicated for the management of (1) mild to moderate pain in adult and pediatric patients 2 years and older; (2) the management of moderate to severe pain with adjunctive opioid analgesics in adult and pediatric patients 2 years and older and (3) the reduction of fever in adult and pediatric patients. **DOSAGE AND ADMINISTRATION:** Adults and adolescents weighing 50 kg and over: the recommended dosage of AMNEPARA DUO™ is 1000 mg every 6 hours or 650 mg every 4 hours, with a maximum single dose of AMNEPARA DUO™ of 1000 mg, a minimum dosing interval of 4 hours, and a maximum daily dose of paracetamol of 4000 mg per day. Adults and adolescents weighing under 50 kg: the recommended dosage of AMNEPARA DUO™ is 15 mg/kg every 6 hours or 12.5 mg/kg every 4 hours, with a maximum single dose of AMNEPARA DUO™ of 15 mg/kg, a minimum dosing interval of 4 hours, and a maximum daily dose of paracetamol of 75 mg/kg per day. Children 2 to 12 years of age: the recommended dosage of AMNEPARA DUO™ is 15 mg/kg every 6 hours or 12.5 mg/kg every 4 hours, with a maximum single dose of AMNEPARA DUO™ of 15 mg/kg, a minimum dosing interval of 4 hours, and a maximum daily dose of paracetamol of 75 mg/kg per day. Neonates, including premature neonates born at ≥ 32 weeks gestational age, up to 28 days chronological age: the recommended dosage of AMNEPARA DUO™ is 12.5 mg/kg every 6 hours, to a maximum daily dose of paracetamol of 50 mg/kg per day, with a minimum dosing interval of 6 hours. Infants 29 days to 2 years of age: the recommended dosage of AMNEPARA DUO™ is 15 mg/kg every 6 hours, to a maximum daily dose of paracetamol of 60 mg/kg per day, with a minimum dosing interval of 6 hours. **ADVERSE REACTIONS:** Hepatic injury, serious skin reactions, allergy and hypersensitivity, anemia, diarrhea, pyrexia, injection site pain, hypokalemia, hypomagnesemia, hypoalbuminemia, hypophosphatemia, muscle spasm, headache, agitation, oliguria, atelectasis, pleural effusion, pulmonary edema, stridor, wheezing, hypotension, and hypertension. **PRECAUTIONS:** Take care when prescribing, preparing, and administering AMNEPARA DUO™ (paracetamol) Injection in order to avoid dosing errors which could result in accidental overdose and death. In particular, be careful to ensure that: (1) the dose in milligrams (mg) and milliliters (ml) is not confused; (2) the dosing is based on weight for patients under 50 kg; (3) infusion pumps are properly programmed; and (4) the total daily dose of paracetamol from all sources does not exceed maximum daily limits. **STORAGE:** Store below 30°C. Do not Freeze. Protect from light. Keep all medicines out of reach of child Discard Unused Portion. **PRESENTATION:** AMNEPARA DUO™ is available in 100 ml in Single-dose Plastic Container. To report SUSPECTED ADVERSE REACTIONS, contact Amneal Healthcare Pvt. Ltd. at 1-800-202-2933 between 9:00 AM and 6:00 PM or visit pv.india@amneal.com.



Amneal Healthcare Private Limited

Regd. Office: 9th Floor, 901 To 905 Icon Elegance, Circle P, S.G. Highway, Ahmedabad 380015

Corporate office: 15-16, Art Guild House, Phoenix Market City, Kurla, Mumbai - 400070

www.amneal.com